

Vanadate

Instructions for use. Sampling kit 2 x 10µl.

6 Capitainer®

INCLUDED IN THE SAMPLING KIT.

Single use products.



Capitainer®B Vanadate sampling card



Disinfectant pad (x2)



lancet (x2)



Disposable Non-adhesive Plaster (x2)



Capitainer®

drying pouch

Additional items you will need:

- A sample ID provided by your kit provider
- Instructions for how to return your sample to the laboratory

PREPARE



- Take a shower
- Exercise
- Hold a cup with hot drink for 1-2 mins
- Soak your hand in warm water



- Read entire instruction before vou start
- Scan QR code or enter https://capitainer.co m/sampling-videob10-vanadate/ to access the video instruction



- Choose a place where you can stand comfortably preferably a dining table. If you prefer to sit, a coffee table works well
- Place all material on table
- Open all packages



- · Open lid of sampling card (a)
- Place card on table
- Do not touch sample wells on card
- Choose a hand to sample from



- Warm your hand for 2-3 mins using a method of choice
- Wash hands with soap and warm water. Dry fully with a clean towel

Cold hands cause unsuccessful sampling!



- Swing your arms to increase blood circulation
- During sampling, keep hand positioned below your elbow



Continue on page two

O2 SAMPLE




Warnings and cautions

- Do not touch the sample inlets before sampling
- Use the card within 30 minutes of opening the package
- Do not bend or twist the card



- Choose a finger to prick, middle or ring finger
- Disinfect with disinfectant pad (b) and allow to dry for 15 seconds



- Twist off and discard lancet (c) cap. DO NOT PULL
- Gently press with your thumb against choosen fingertip to fill it with blood



- Place lancet (c) against side of fingertip and press firmly
- Lancet (c) activates automatically and retracts into device



- Release pressure on fingertip and wait for the blood to appear
- Wipe off first drop with the nonadhesive pad (d)
- Let a new large drop of blood form by itself (wait for the drop to almost fall off by its own weight)
- Apply it to one of the sample wells

Apply only one drop. More drops may result in an incorrect sample.


5 Important step, read carefully!



- Let a new large drop of blood form by itself (wait for the drop to almost fall off by its own weight)
- Apply it to the **second** sample well

Apply only one drop. More drops may result in an incorrect sample.

6



- Stop bleeding with non-adhesive pad
 (d)
- Apply plaster (e)
- Check for sampling success: upon successful sampling, the control windows should be showing red as shown in picture

03

PACK & RETURN

All used materials such as cards, lancets, gauzes, swabs with blood residues must be handled and disposed of safely and in accordance with local regulations.



- · Label your sampling card
- Place sampling card with lid closed in drying pouch (f)
- Return sample as directed by your test provider

■ Capitainer®Rev. Date 2025-11-13